Sub-national estimation of population size for female sex workers in South Africa: Review and Bayesian evidence synthesis

### Nanina Anderegg<sup>1,2</sup>, Leigh F Johnson<sup>2</sup>

<sup>1</sup>Institute of Social and Preventive Medicine (ISPM), University of Bern, Switzerland

<sup>2</sup>Centre for Integrated Data and Epidemiological Research (CIDER), Unversity of Cape Town, South Africa



How many female sex workers (FSW) are there in South Africa?





How many female sex workers (FSW) are there in the North West Province?





How many female sex workers (FSW) are there in the Central Karoo district?





How many female sex workers (FSW) are there in the Alfred Duma sub-district?







• high-risk/key population for HIV



### high-risk/key population for HIV

• tailored interventions, resource allocation, equitable & equal service access



### high-risk/key population for HIV

• tailored interventions, resource allocation, equitable & equal service access



### high-risk/key population for HIV

• tailored interventions, resource allocation, equitable & equal service access

hidden population



### high-risk/key population for HIV

• tailored interventions, resource allocation, equitable & equal service access

#### hidden population

criminalization and stigmatization



### high-risk/key population for HIV

• tailored interventions, resource allocation, equitable & equal service access

#### hidden population

criminalization and stigmatization



### high-risk/key population for HIV

• tailored interventions, resource allocation, equitable & equal service access

hidden population

criminalization and stigmatization

Some studies reported population size estimates (PSE) for FSW



### high-risk/key population for HIV

• tailored interventions, resource allocation, equitable & equal service access

hidden population

criminalization and stigmatization

Some studies reported population size estimates (PSE) for FSW

Often estimate for a small-area level + extrapolation



### high-risk/key population for HIV

• tailored interventions, resource allocation, equitable & equal service access

3/12

#### hidden population

criminalization and stigmatization

- Some studies reported population size estimates (PSE) for FSW
- Often estimate for a small-area level + extrapolation
- Differences in estimation method, catchment area, year ...

### high-risk/key population for HIV

• tailored interventions, resource allocation, equitable & equal service access

### hidden population

criminalization and stigmatization

- Some studies reported population size estimates (PSE) for FSW
- Often estimate for a small-area level + extrapolation
- Differences in estimation method, catchment area, year ...
- Lack of "synthesized" PSEs on sub-district and district level



## objective

To estimate population size of FSW in South Africa on sub-national level by leveraging data from different existing data sources



# methods

published PSE for FSW on sub-district or lower level
demographic data on South African sub-districts



## methods

published PSE for FSW on sub-district or lower level
demographic data on South African sub-districts

Bayesian hierarchical regression model

- outcome: proportion FSW among females aged 15-64
- fixed effects: province, boarder, pop density, urbanization index (+ interaction)
- random effects & weights (to account for dependence & estimation method)



## methods

published PSE for FSW on sub-district or lower level
demographic data on South African sub-districts

Bayesian hierarchical regression model

- outcome: proportion FSW among females aged 15-64
- fixed effects: province, boarder, pop density, urbanization index (+ interaction)
- random effects & weights (to account for dependence & estimation method)

% FSW at sub-district level (extrapolation to areas w/o. published PSE).
aggregate predictions for district-, province-, national-level estimates





### Published FSW proportions.





#### Published FSW proportions.





5 studies (2013-2018)

- 115 published PSE (including extrapolated, duplicated, combination and implausible)
- 74 published PSE included: 1.2%-5.8%

#### Published FSW proportions.





 published PSE for 45 (out of 213) sub-districts in 31 (out of 52) districts





- published PSE for 45 (out of 213) sub-districts in 31 (out of 52) districts
- mostly 1 published PSE, but for some up to 10







Predicted sub-district % FSW.







Predicted sub-district % FSW.





max: 5.4% (3.1-10.9) min: 0.2% (0.1-0.4) Proportion FSW [%] 1 2 3 4 5

Predicted sub-district % FSW.















9/12





 large variability in predicted % FSW on sub-district level, reduced on district and province level



- large variability in predicted % FSW on sub-district level, reduced on district and province level
- quality of input data (reported PSE)?



- large variability in predicted % FSW on sub-district level, reduced on district and province level
- quality of input data (reported PSE)?
- high uncertainty in estimates



- large variability in predicted % FSW on sub-district level, reduced on district and province level
- quality of input data (reported PSE)?
- high uncertainty in estimates
- outlook: include routine data (PrEP), spatial correlation



How many female sex workers (FSW) are there in South Africa?

0.9% (0.6-1.6%) among females aged 15-64





How many female sex workers (FSW) are there in the North West Province?

0.5% (0.1-2.0%) among females aged 15-64





How many female sex workers (FSW) are there in the Central Karoo district?

2.1% (0.9-5.3%) among females aged 15-64





How many female sex workers (FSW) are there in the Alfred Duma sub-district?

1.6% (0.6-4.7%) among females aged 15-64





### acknowledgements

- Leigh Johnson
- Oliver Stevens

#### Swiss National Science Foundation

Nanina Anderegg was supported by funding from the Swiss National Science Foundation (grant no. P500PM\_203010, P5R5PM\_225275) and the International epidemiology Databases to Evaluate AIDS (IeDEA) collaboration.



The International Epidemiology Databases to Evaluate AIDS (IeDEA) is supported by the U.S. National Institutes of Health's National Institute of Allergy and Infectious Diseases, the Eunice Kennedy Shriver National Institute of Child Health and Human Development, the National Cancer Institute, the National Institute of Mental Health, the National Institute on Drug Abuse, the National Heart, Lung, and Blood Institute, the National Institute on Alcohol Abuse and Alcoholism, the National Institute of Diabetes and Digestive and Kidney Diseases, and the Fogarty International Center: Asia-Pacific, U01AI069907; CCASAnet, U01AI069923; Central Africa, U01AI096299; East Africa, U01AI069911; NA-ACCORD, U01AI069918; Southern Africa, U01AI069924; West Africa, U01AI069919. Informatics resources are supported by the Harmonist project, R24AI24872. This work is solely the responsibility of the authors and does not necessarily represent the official views of any of the institutions mentioned above.

